

Exterior splashes and linear sets of rank 3

S.G. Barwick and Wen-Ai Jackson

20 April 2014

Corresponding Author: Dr Susan Barwick, University of Adelaide, Adelaide 5005, Australia.
Phone: +61 8 8313 3983, Fax: +61 8 8313 3696, email: susan.barwick@adelaide.edu.au

Keywords: subplanes, sublines, linear sets, Sherk surfaces, circle geometries

AMS code: 51E20

Abstract

In $\text{PG}(2, q^3)$, let π be a subplane of order q that is exterior to ℓ_∞ . The exterior splash of π is defined to be the set of $q^2 + q + 1$ points on ℓ_∞ that lie on a line of π . This article investigates properties of an exterior order- q -subplane and its exterior splash. We show that the following objects are projectively equivalent: exterior splashes, covers of the circle geometry $CG(3, q)$, Sherk surfaces of size $q^2 + q + 1$, and scattered linear sets of rank 3. We compare our construction of exterior splashes with the projection construction of a linear set. We give a geometric construction of the two different families of sublines in an exterior splash, and compare them to the known families of sublines in a scattered linear set of rank 3.

1 Introduction

Let π be a subplane of $\text{PG}(2, q^3)$ of order q that meets ℓ_∞ in 0 or 1 point. Each line of π , when extended to $\text{PG}(2, q^3)$, meets ℓ_∞ in a point. The set of points of ℓ_∞ that lie on a line of π is called the *splash* of π . If π is tangent to ℓ_∞ at a point T , then the splash of π consists of $T = \pi \cap \ell_\infty$, and q^2 further points. We call this splash a *tangent splash*, and these were investigated in [3, 4]. If π is exterior to ℓ_∞ , then the splash $\mathbb{S}$ contains $q^2 + q + 1$ points, and we call it an *exterior splash*. Note that each line of π contains a distinct point of $\mathbb{S}$, and conversely, each point of $\mathbb{S}$ lies on a unique line of π . More generally, we can define an exterior splash onto any exterior line. That is, let π be a subplane of order q of $\text{PG}(2, q^3)$, and let ℓ be an exterior line of π , then the *exterior splash of π onto ℓ* is the set of $q^2 + q + 1$ points of ℓ that lie on lines of π .

In this article we will investigate exterior order- q -subplanes and exterior splashes. In Section 3 we consider the relationship of exterior splashes to other known objects. In particular, exterior splashes are projectively equivalent to scattered linear sets of rank 3; covers of circle geometries $CG(3, q)$; and Sherk surfaces of size $q^2 + q + 1$.

The rest of the article focuses on the relationship between an exterior order- q -subplane and its exterior splash. In Section 4, we use order- q -sublines contained in an exterior splash $\mathbb{S}$ in terms of an associated order- q -subplane π . In an analogous manner to [12, Remark 20], we also discuss why projective bundles of conics of π arise naturally as order- q -sublines of $\mathbb{S}$. In Section 4.2, we show that the two families of $q^2 + q + 1$ sublines in an exterior splash are equivalent in some sense, despite their apparent difference in definition.

A $\text{GF}(q)$ -linear set of rank 3 can be obtained by projecting an order- q -subplane of $\text{PG}(2, q^3)$ onto a line, see [16]. In Section 5.1, we investigate the relationship between this projection construction of a scattered linear set of rank 3, and our construction of an exterior splash. In Section 5.2, we revisit the tangent splash of a tangent order- q -subplane, and look at the projection in this context.

Finally, in Section 6, we look at two exterior order- q -subplanes that have a common exterior splash, and determine how they can intersect. Further, we show that there are exactly two order- q -subplanes that have a common exterior splash and share a fixed order- q -subline.

2 Notation and definitions

In this section we introduce the notation we use, as well as defining linear sets, circle geometries and Sherk surfaces.

If π is a subplane of $\text{PG}(2, q^3)$ of order q , then we call π an *order- q -subplane*. If ℓ_∞ is an exterior line of π , then we say π is an *exterior order- q -subplane*. An *order- q -subline* of $\text{PG}(2, q^3)$ is a line of an order- q -subplane, that is, it is isomorphic to $\text{PG}(1, q)$. Points in $\text{PG}(2, q^3)$ have homogeneous coordinates (x, y, z) with $x, y, z \in \text{GF}(q^3)$. Let the line at infinity ℓ_∞ have equation $z = 0$; so the affine points of $\text{PG}(2, q^3)$ have coordinates $(x, y, 1)$. We can construct $\text{GF}(q^3)$ as a cubic extension of $\text{GF}(q)$ using a primitive element τ with primitive polynomial

$$x^3 - t_2x^2 - t_1x - t_0, \tag{1}$$

where $t_0, t_1, t_2 \in \text{GF}(q)$; so every element in $\text{GF}(q^3)$ can be uniquely written as $a_0 + a_1\tau + a_2\tau^2$ with $a_0, a_1, a_2 \in \text{GF}(q)$.

We define linear sets of $\text{PG}(1, q^3)$, more generally, linear sets of $\text{PG}(n - 1, q^t)$ are defined in [12]. The points of $\text{PG}(1, q^3)$ can be considered as elements of the 2-dimensional vector space $V = \text{GF}(q^3)^2$ over $\text{GF}(q^3)$. Let U be a subset of V that forms a 3-dimensional vector space over $\text{GF}(q)$. Then the vectors of U (considered as vectors over $\text{GF}(q^3)$) form a $\text{GF}(q)$ -linear set of rank 3 of $\text{PG}(1, q^3)$. A *scattered linear set of rank 3* is a $\text{GF}(q)$ -linear set of $\text{PG}(1, q^3)$ of

rank 3 and size $q^2 + q + 1$. By [12], all scattered linear sets of rank 3 are projectively equivalent. Scattered linear sets were introduced in [5], and have recently been studied in [11, 12, 13, 15].

In [8, 9], Bruck gave a set of axioms for higher dimensional circle geometries. We look at the 3-dimensional case, namely $CG(3, q)$. The points of $CG(3, q)$ can be identified with the points of $PG(1, q^3)$, and the *circles* of $CG(3, q)$ are identified with the order- q -sublines of $PG(1, q^3)$. The *stability group* of a circle is defined to be the subgroup of $\text{Aut}CG(3, q) = \text{P}\Gamma L(2, q^3)$ that fixes the circle pointwise. For two distinct points of $CG(3, q)$, let $\phi(P, Q)$ denote the group generated by the stability groups of all circles containing P and Q . A *cover* is defined to be an orbit under $\phi(P, Q)$ of any point R distinct from P, Q . In [9], it is shown that every cover of $CG(3, q)$ has $q^2 + q + 1$ points, and can be represented in one of the following two ways, using the identification of the points of $CG(3, q)$ with the field $\text{GF}(q^3) \cup \{\infty\}$:

- I. $\{x \in \text{GF}(q^3) : N(x - a) = f\}$, for some $a \in \text{GF}(q^3)$, and $f \in \text{GF}(q) \setminus \{0\}$.
- II. $\{x \in \text{GF}(q^3) \cup \{\infty\} : N\left(\frac{x-a}{x-b}\right) = f\}$, for some $a, b \in \text{GF}(q^3)$, and $f \in \text{GF}(q) \setminus \{0\}$.

where N is the norm from $\text{GF}(q^3)$ to $\text{GF}(q)$, that is $N(x) = x^{q^2+q+1}$. The points P, Q are called the *carriers* of the cover. The carriers for a cover of type I are $\{a, \infty\}$, and the carriers for a cover of type II are $\{a, b\}$. A cover of type II will contain ∞ if and only if $f = 1$.

Sherk [17], studied objects other than circles and covers in $CG(3, q)$. Representing points of $CG(3, q)$ as elements of $\text{GF}(q^3) \cup \{\infty\}$, a *Sherk surface* is the set of points satisfying

$$S(f, \alpha, \delta, g) = \{z \in \text{GF}(q^3) \cup \{\infty\} : fN(z) + T(\alpha^{q^2}z^{q+1}) + T(\delta z) + g = 0\}$$

for $f, g \in \text{GF}(q)$, $\alpha, \delta \in \text{GF}(q^3)$, where N, T are the norm and trace from $\text{GF}(q^3)$ to $\text{GF}(q)$. Sherk surfaces of size $q^2 + q + 1$ are precisely the Bruck covers of $CG(3, q)$.

3 Exterior splashes

3.1 An important Singer group

We will need a particular Singer group acting on order- q -subplanes. First we define the notion in $PG(2, q^3)$ of conjugate points and lines with respect to an order- q -subplane. Let π be an order- q -subplane of $PG(2, q^3)$. There is a unique collineation group of order three that fixes every point of π , let ζ be a generator of this group. Then the points of $PG(2, q^3) \setminus \pi$ can be partitioned into sets of size three of form $\{P, \zeta(P), \zeta^2(P)\}$, called *conjugate points with respect to π* . Note that $\{P, \zeta(P), \zeta^2(P)\}$ are collinear if and only if they lie on the extension of an order- q -subline of π . Similarly, the lines of $PG(2, q^3) \setminus \pi$ can be partitioned into sets of size three of form $\ell, \zeta(\ell), \zeta^2(\ell)$, called *conjugate lines with respect to π* . For example, let $\pi = PG(2, q)$, then $\zeta : (x, y, z) \mapsto (x^q, y^q, z^q)$ fixes every point of π and acts semi-regularly on the remaining points of $PG(2, q^3)$. If P is a point of $PG(2, q^3) \setminus PG(2, q)$, then the conjugate points with respect to the order- q -subplane $\pi = PG(2, q)$ are P, P^q, P^{q^2} . If ℓ is a line of $PG(2, q^3) \setminus PG(2, q)$, then the conjugate lines with respect to the order- q -subplane $\pi = PG(2, q)$ are ℓ, ℓ^q, ℓ^{q^2} .

We need the following result about the collineation groups acting on an exterior order- q -subplane.

Theorem 3.1 *Consider the collineation group $G = \text{PGL}(3, q^3)$ acting on $\text{PG}(2, q^3)$. Let $I = G_{\pi, \ell}$ be the subgroup of G fixing an order- q -subplane π , and a line ℓ exterior to π . Then*

1. *I is cyclic of order $q^2 + q + 1$, and acts regularly on the points and on the lines of π .*
 2. *I fixes exactly three lines: ℓ , and its conjugates m, n with respect to π . Further I acts semi-regularly on the remaining line orbits.*
 3. *I fixes exactly three points: $E_1 = \ell \cap m, E_2 = \ell \cap n, E_3 = m \cap n$ (which are conjugate with respect to π) and acts semi-regularly on the remaining point orbits.*
- The points E_1, E_2 are called the carriers of π .*

Proof We first consider the homography ϕ of $\text{PG}(2, q^3)$ with matrix

$$T = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ t_0 & t_1 & t_2 \end{pmatrix},$$

where t_0, t_1, t_2 are as in (1) and $\phi(x, y, z) = T(x, y, z)^t$. The matrix T has three eigenvalues τ, τ^q and τ^{q^2} with corresponding eigenvectors $Q = (1, \tau, \tau^2), Q^q$ and Q^{q^2} respectively. Further, the eigenvalues of T^i are $\tau^i, \tau^{iq}, \tau^{iq^2}$ with corresponding eigenvectors Q, Q^q, Q^{q^2} respectively. Now $\tau^i = \tau^{iq}$ if and only if $\tau^i \in \text{GF}(q)$, if and only if $q^2 + q + 1 \mid i$, hence the eigenvalues are distinct. Thus, for $0 < i < q^2 + q + 1$, ϕ^i fixes exactly three points Q, Q^q, Q^{q^2} . Further, ϕ has order $q^2 + q + 1$, and ϕ acts semi-regularly on the points other than Q, Q^q, Q^{q^2} . In particular, $\langle \phi \rangle$ is a Singer cycle of the order- q -subplane $\pi = \text{PG}(2, q)$, and so acts regularly on the points and on the lines of π .

Now let $G = \text{PGL}(3, q^3)$, it is well known that $|G| = q^9(q^9 - 1)(q^6 - 1)$, and that G is transitive on the points of $\text{PG}(2, q^3)$; on the lines of $\text{PG}(2, q^3)$; and on the order- q -subplanes of $\text{PG}(2, q^3)$. Hence when considering the subgroup $K = G_\pi$ for some order- q -subplane π , we can without loss of generality let $\pi = \text{PG}(2, q)$. Let $K = G_\pi = \text{PGL}(2, q)$. Thus K is transitive on the points and lines of π , and so this gives one point orbit of K . Further, $|K| = q^3(q^3 - 1)(q^2 - 1)$.

We now consider the point $P = (1, 1, \tau)$ and calculate its orbit under K . Note that P is not in π , but P is on the line $[1, -1, 0]$, which is a line of π , so P lies on exactly one line of π . Consider the subgroup of K fixing P , so consider a homography with matrix

$$C = \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix},$$

with $a, \dots, i \in \text{GF}(q)$, such that $C(1, 1, \tau)^t \equiv (1, 1, \tau)$. Then $a + b + c\tau = d + e + f\tau$ (hence $c = f$ and $a + b = d + e$) and $(a + b + c\tau)\tau = g + h + i\tau$ giving $c = 0$ (so $f = 0$), $a + b = i$, and $g + h = 0$. Hence

$$C = \begin{pmatrix} a & b & 0 \\ d & a + b - d & 0 \\ g & -g & a + b \end{pmatrix} = \begin{pmatrix} a & 1 - a & 0 \\ d & 1 - d & 0 \\ g & -g & 1 \end{pmatrix}$$

since we can take $a + b = 1$ as $\det C \neq 0$. Now $\det C \neq 0$ if and only if $a \neq d$. Hence the size of K_P is $q^2(q - 1)$. Thus by the orbit stabilizer theorem, $|P^K| = |K|/|K_P| = q(q^3 - 1)(q + 1)$. This is the number of points of $\text{PG}(2, q^3) \setminus \pi$ that lie on exactly one line of π , hence these points lie in one orbit of K .

We now consider the points of $\text{PG}(2, q^3) \setminus \pi$ that lie on exactly zero lines of π . Consider the point $Q = (1, \tau, \tau^2)$, it does not lie on any line of $\text{PG}(2, q)$. We calculate the size of the orbit of Q under K . Note first that the subgroup of K generated by the homography ϕ with matrix T fixes Q . Now consider a homography of K with matrix

$$B = \begin{pmatrix} a & b & c \\ d & e & f \\ g & h & i \end{pmatrix},$$

$a, \dots, i \in \text{GF}(q)$ that fixes Q . We have $(a + b\tau + c\tau^2)\tau = d + e\tau + f\tau^2$ and $(d + e\tau + f\tau^2)\tau = g + h\tau + i\tau^2$. So we have $B = (\mathbf{x}, \mathbf{x}T, \mathbf{x}T^2)^t$ where $\mathbf{x} = (a, b, c)$. Now B has nonzero determinant if and only not all of a, b, c are zero. Further, $\mathbf{x} = (a, b, c)$ and $\mathbf{x} = y(a, b, c)$ correspond to the same collineation if and only if $y \in \text{GF}(q)$. Thus the subgroup K_Q is of size $q^2 + q + 1$. Hence K_Q is generated by the Singer cycle ϕ with matrix T . By the orbit stabilizer theorem, $|Q^K| = |K|/|K_Q| = q^3(q^2 - 1)(q - 1)$. This is equal to the number of points of $\text{PG}(2, q^3)$ that lie on zero lines of $\pi = \text{PG}(2, q)$. Hence these points all lie in one orbit of K . A similar argument shows that K has three line orbits: the lines of π , the lines meeting π in exactly one point, and the lines exterior to π .

As $K = G_\pi$ is transitive on the exterior lines of π , so we can without loss of generality consider the line $\ell = [1, \tau, \tau^2]$ which is exterior to $\pi = \text{PG}(2, q)$, and we calculate $I = G_{\pi, \ell}$ for this π, ℓ . Consider the homography ϕ' which has matrix T^{-t} (the transpose of the inverse of T). Note that ϕ' has order $q^2 + q + 1$ as ϕ does. Now ϕ' maps a line $[l, m, n]$ of $\text{PG}(2, q^3)$ to the line $[l, m, n]T^t = \ell T^t$. Hence ϕ' fixes the line $\ell = [1, \tau, \tau^2]$. So using a similar argument to the above paragraph, we have $I = G_{\pi, \ell}$ is generated by the singer cycle ϕ' . Hence $|I| = q^2 + q + 1$ and I acts regularly on the points, and on the lines, of $\text{PG}(2, q)$, proving part 1.

As T has entries in $\text{GF}(q)$, we have $T = T^q$, hence $\phi'(\ell^q) = \ell^q T^t = (\ell T^t)^q = (\ell)^q = \ell^q$. That is ϕ' fixes ℓ^q , similarly, ϕ' fixes ℓ^{q^2} . Hence ϕ' will fix the three points $P = \ell \cap \ell^{q^2}$, $P^q = \ell \cap \ell^q$, $P^{q^2} = \ell^q \cap \ell^{q^2}$. A similar argument to that used when analysing the homography ϕ shows that I acts semi-regularly on the remaining line (and point) orbits of $\text{PG}(2, q^3)$, proving parts 2 and 3. $\square$

The Singer group $I = \text{PGL}(3, q^3)_{\pi, \ell}$ has a number of important consequences. We can use I to define a special family of conics and dual conics in π which play an important role in exterior splashes. A conic of π whose extension to $\text{PG}(2, q^3)$ contains the three conjugate points E_1, E_2, E_3 is called a (π, ℓ) -special conic of π . Dually we can define a (π, ℓ) -special-dual conic to be an dual conic of π that contains the three conjugate lines ℓ, m, n . Special conics and dual conics are irreducible. Further, the two incidence structures with points the points of π , lines the (π, ℓ) -special conics (respectively dual conics) of π , and natural incidence are isomorphic to $\text{PG}(2, q)$. In addition, a *projective bundle* of conics is defined in [1] to be a set of $q^2 + q + 1$ conics of $\text{PG}(2, q)$ which pairwise meet in exactly one point. There are only three

known classes of projective bundles, of which the (π, ℓ) -special conics of π form a *circumscribed bundle*, and the (π, ℓ) -special-dual conics of π form an *inscribed bundle*.

3.2 An example

We will need an example of an order- q -subplane to prove subsequent results. So the next result takes the order- q -subplane $\pi = \text{PG}(2, q)$ of $\text{PG}(2, q^3)$, finds a suitable exterior line ℓ to π , and calculates the exterior splash, the Singer group I , and the carriers for π .

Example 3.2 Let $\pi = \text{PG}(2, q)$ be an order- q -subplane of $\text{PG}(2, q^3)$.

1. The line $\ell = [-\tau\tau^q, \tau^q + \tau, -1]$ is exterior to π .
2. The exterior splash $\mathbb{S}$ of $\pi = \text{PG}(2, q)$ on $\ell = [-\tau\tau^q, \tau^q + \tau, -1]$ is

$$\mathbb{S} = \{E + \theta E^q : \theta^{q^2+q+1} = -1, \theta \in \text{GF}(q^3)\}, \quad \text{where } E = (1, \tau, \tau^2).$$

3. The Singer group $I = \text{PGL}(3, q^3)_{\pi, \ell}$ is generated by the homography ϕ with matrix

$$T = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ t_0 & t_1 & t_2 \end{pmatrix},$$

where t_0, t_1, t_2 are as in (1), and $\phi(x, y, z)^t = T(x, y, z)^t$. The three fixed points of ϕ are the points $E = \ell^{q^2} \cap \ell = (1, \tau, \tau^2)$, E^q, E^{q^2} . The three fixed lines of ϕ are ℓ, ℓ^q, ℓ^{q^2} . So π has carriers $E = (1, \tau, \tau^2)$ and $E^q = (1, \tau^q, \tau^{2q})$.

Proof We first show that the line $\ell = [-\tau\tau^q, \tau^q + \tau, -1]$ is exterior to $\pi = \text{PG}(2, q)$. The points of π are of the form (x, y, z) , $x, y, z \in \text{GF}(q)$, not all zero. This point lies on the line $\ell = [-\tau\tau^q, \tau^q + \tau, -1]$ if and only if $\tau\tau^q x - (\tau + \tau^q)y + z = 0$ for some $x, y, z \in \text{GF}(q)$, not all zero. As $1, \tau\tau^q, \tau + \tau^q$ of $\text{GF}(q^3)$ are linearly independent over $\text{GF}(q)$, $\tau\tau^q x - (\tau + \tau^q)y + z = 0$ has no solutions with $x, y, z \in \text{GF}(q)$, not all zero, thus π is exterior to ℓ . We can parameterize the line ℓ as $\ell = \{E + \theta E^q : \theta \in \text{GF}(q^3) \cup \{\infty\}\}$, where $E = (1, \tau, \tau^2)$ (so E has parameter 0, and E^q has parameter ∞). Consider a line ℓ_i of π , so $\ell_i = [l, m, n]$, $l, m, n \in \text{GF}(q)$, not all zero. Then ℓ_i meets the line $\ell = [-\tau\tau^q, \tau^q + \tau, -1]$ in the point $X_i = (m + n(\tau + \tau^q), n\tau\tau^q - l, -l(\tau + \tau^q) - m\tau\tau^q)$. We can write this as $X_i = E + \theta_i E^q$ where $\theta_i = -(l + m\tau + n\tau^2)/(l + m\tau + n\tau^2)^q$. Note that $\theta_i^{q^2+q+1} = \theta_i^{q^2} \theta_i^q \theta_i = -1$. As there are $q^2 + q + 1$ solutions to the equation $\theta^{q^2+q+1} = -1$, the $q^2 + q + 1$ lines of π meet ℓ in the $q^2 + q + 1$ points $E + \theta E^q$ with $\theta^{q^2+q+1} = -1$. That is, π has exterior splash $\mathbb{S} = \{E + \theta E^q : \theta^{q^2+q+1} = -1, \theta \in \text{GF}(q^3)\}$ on ℓ , completing the proof of part 2. The proof of part 3 involves straightforward calculations. $\square$

3.3 Exterior splashes and linear sets

The following result is important to our study of exterior order- q -subplanes and exterior splashes.

Theorem 3.3 *Consider the collineation group $G = \text{PGL}(3, q^3)$ acting on $\text{PG}(2, q^3)$. The subgroup G_ℓ fixing a line ℓ is transitive on the order- q -subplanes that are exterior to ℓ , and is transitive on the exterior splashes on ℓ .*

Proof Recall that the group $G = \text{PGL}(3, q^3)$ is transitive on the order- q -subplanes of $\text{PG}(2, q^3)$. From the proof of Theorem 3.1, the subgroup of K fixing an order- q -subplane π is transitive on the exterior lines of π . Hence G is transitive on pairs (π, ℓ) where π is an order- q -subplane and ℓ is a line exterior to π . Thus G_ℓ is transitive on the order- q -subplanes that are exterior to ℓ , and hence G_ℓ is transitive on the exterior splashes of ℓ . $\square$

The importance of this result is two fold. Firstly, it means that when proving results about exterior order- q -subplanes and exterior splashes, we can without loss of generality prove results for a particular example order- q -subplane. Secondly, this theorem shows that all exterior splashes are projectively equivalent. It is then straightforward to use Example 3.2, and the result that all scattered linear sets of rank 3 are projectively equivalent by [12], to prove that exterior splashes of $\text{PG}(2, q^3)$ are projectively equivalent to scattered linear sets of rank 3.

Theorem 3.4 *Exterior splashes of $\text{PG}(1, q^3)$ are projectively equivalent to scattered linear sets of rank 3.*

We note that in concurrent, independent work, Lavrauw and Zanella [14] prove the more general result that exterior splashes of $\text{PG}(1, q^n)$ are projectively equivalent to scattered linear sets of $\text{PG}(1, q^n)$.

3.4 Circle geometries and Sherk surfaces

In this section we observe that exterior splashes are projectively equivalent to covers of the circle geometry $CG(3, q)$, and to Sherk surfaces of size $q^2 + q + 1$.

Theorem 3.5 *Let π be an exterior order- q -subplane with exterior splash $\mathbb{S}$ on $\ell_\infty = \text{PG}(1, q^3)$ and carriers E_1, E_2 . Then $\mathbb{S}$ is projectively equivalent to a cover of the circle geometry $CG(3, q)$ with carriers E_1, E_2 .*

Proof By Theorem 3.3, we can without loss of generality prove this for the exterior splash $\mathbb{S}$ of the order- q -subplane $\pi = \text{PG}(2, q)$ onto the exterior line $\ell = [-\tau\tau^q, \tau^q + \tau, -1]$. By Example 3.2, $\mathbb{S}$ has exterior splash $\mathbb{S} = \{E + \theta E^q : \theta^{q^2+q+1} = -1, \theta \in \text{GF}(q^3)\}$ on ℓ , and carriers $E_1 = E = (1, \tau, \tau^2)$ and $E_2 = E^q$ (corresponding to $\theta = 0$ and $\theta = \infty$ respectively). Hence using the notation of Section 2, the exterior splash is a cover of type I with $a = 0$, $f = -1$. $\square$

We note that this equivalence can also be deduced from the general theory of linear sets of pseudoregulus type in [15]. In particular, an exterior splash $\mathbb{S}$ is equivalent to a scattered linear set of rank 3 of pseudoregulus type. The carriers of $\mathbb{S}$ are called transversal spaces of the pseudoregulus. There are two important corollaries for exterior splashes, we note that they both follow immediately from properties of circle geometries, and are also proved directly in [15]. We state them here in the exterior splash context as they are important for understanding the carriers of an exterior splash.

Result 3.6 *1. Let π_1, π_2 be two exterior order- q -subplanes with the same exterior splash $\mathbb{S}$ on ℓ_∞ , then π_1 and π_2 have the same carriers. That is, an exterior splash has two unique carriers.*
2. Given two carriers E_1, E_2 of ℓ_∞ , there are exactly $q - 1$ disjoint exterior splashes with carriers E_1, E_2 .

Remark 3.7 As the Sherk surfaces of size $q^2 + q + 1$ are precisely the Bruck covers of $CG(3, q)$, exterior splashes are also projectively equivalent to Sherk surfaces of size $q^2 + q + 1$. In fact, the Sherk surface corresponding to the exterior splash of the order- q -subplane $\pi = \text{PG}(2, q)$ onto the exterior line $\ell = [-\tau\tau^q, \tau^q + \tau, -1]$ has parameters $S(1, 0, 0, 1)$. This equivalence means we can adapt results about permutation groups of Sherk surfaces from [17] to our setting, also see [11]. In particular we will need the following collineation group.

Lemma 3.8 *Let $\mathbb{S}$ be the exterior splash of the order- q -subplane $\pi = \text{PG}(2, q)$ onto the exterior line $\ell = [-\tau\tau^q, \tau^q + \tau, -1]$. Then $\mathbb{S} \equiv \{(\theta, 1) : \theta^{q^2+q+1} = -1, \theta \in \text{GF}(q^3)\}$, where $(\theta, 1) \equiv \theta E_1 + E_2$. The group of collineations of ℓ fixing $\mathbb{S}$ has as generators two homographies Γ, Δ with matrices*

$$\begin{pmatrix} \tau & 0 \\ 0 & \tau^q \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$$

respectively; so Γ fixes the carriers E_1, E_2 , and Δ interchanges them.

4 Order- q -sublines of an exterior splash

We have so far discussed the relation between exterior splashes and other geometrical objects in their own right. For the remainder of the paper we will primarily consider the relationship between an exterior order- q -subplane and its exterior splash. In this section we investigate the order- q -sublines contained in an exterior splash. We first characterise the order- q -sublines of an exterior splash *with respect to* an associated exterior order- q -subplane. Then we compare this with the characterisation of order- q -sublines of a scattered linear set discussed in [12].

4.1 Sublines in an exterior splash

We first note that the dual of [12, Corollary 19] can be generalised to the following result.

Lemma 4.1 *Let π be an exterior order- q -subplane of $\text{PG}(2, q^3)$ with exterior splash $\mathbb{S}$ on ℓ_∞ , and let Γ be a dual $(q+1)$ -arc of π . The lines of Γ meet the exterior splash $\mathbb{S}$ in an order- q -subline if and only if Γ is (π, ℓ_∞) -special-dual conic of π .*

Using this lemma, it is straightforward to prove the following geometric interpretation of the order- q -sublines of $\mathbb{S}$ in relation to an associated order- q -subplane.

Theorem 4.2 *Let π be an exterior order- q -subplane of $\text{PG}(2, q^3)$, $q > 2$, with exterior splash $\mathbb{S}$ on ℓ_∞ . The $2(q^2 + q + 1)$ order- q -sublines of $\mathbb{S}$ lie in two families of size $q^2 + q + 1$ as follows.*

1. *If A is a point of π , then the pencil of $q+1$ lines of π through A meets ℓ_∞ in an order- q -subline of $\mathbb{S}$, called a π -pencil-subline.*
2. *If Γ is a (π, ℓ_∞) -special-dual conic of π , then the lines of Γ meet ℓ_∞ in an order- q -subline of $\mathbb{S}$, called a π -dual-conic-subline.*

Remark 4.3 The π -dual-conic-sublines of $\mathbb{S}$ arise from an inscribed bundle of conics (see Section 3.1). We note that in [12, Remark 20], the authors show that irregular sublines of a scattered linear set of rank 3 arise from a circumscribed bundle of conics.

4.2 Sublines of a linear set and an exterior splash

In this section we compare the construction of the two families of order- q -sublines from [12] with that of Theorem 4.2. In particular, we explain why the notions of regular and irregular order- q -sublines of a linear set defined in [12] are not intrinsic properties of the linear set, but that the two classes can be interchanged.

We first prove that the characterisation of order- q -sublines given in Theorem 4.2 is a property of the associated order- q -subplane, not a property of the exterior splash.

Theorem 4.4 *Consider an exterior splash $\mathbb{S}$ of an exterior order- q -subplane π onto ℓ_∞ . Let $\mathcal{X}$ denote the π -pencil-sublines of $\mathbb{S}$, and let $\mathcal{Y}$ denote the π -dual-conic-sublines of $\mathbb{S}$. Then there exists an exterior order- q -subplane π' with the same exterior splash $\mathbb{S}$ on ℓ_∞ , such that $\mathcal{X}$ contains the π' -dual-conic-sublines of $\mathbb{S}$ and $\mathcal{Y}$ contains the π' -pencil-sublines of $\mathbb{S}$.*

Proof By Theorem 3.3, all exterior splashes are projectively equivalent, so we can without loss of generality look at the exterior splash $\mathbb{S}$ of Example 3.2. Now $\mathbb{S}$ contains the points $X_i = E + \theta_i E^q$, so points in $\mathbb{S}$ have coordinates equivalent to $(\theta_i, 1) \equiv (l + m\tau + n\tau^2, -(l + m\tau + n\tau^2)^q)$ where $l, m, n \in \text{GF}(q)$, not all zero. We can map this to the point $(l + m\tau + n\tau^2, (l + m\tau + n\tau^2)^q, 0)$, so, we can without loss of generality let $\mathbb{S} = \{(x, x^q, 0) : x \in \text{GF}(q^3)\}$. (We note that this is equivalent to the canonical form for a linear set of pseudoregulus type given in [15].) It

is straightforward to calculate the order- q -sublines of $\mathbb{S}$, they are $\mathcal{X} = \{x_P : P \in \text{PG}(2, q)\}$, $\mathcal{Y} = \{y_P : P \in \text{PG}(2, q)\}$, where

$$\begin{aligned} x_P &= \{(l + m\tau + n\tau^2, (l + m\tau + n\tau^2)^q, 0) : l, m, n \in \text{GF}(q), P \cdot [l, m, n] = 0\} \\ y_P &= \{((l + m\tau + n\tau^2)^q, l + m\tau + n\tau^2, 0) : l, m, n \in \text{GF}(q), P \cdot [l, m, n] = 0\}. \end{aligned}$$

Let π be an exterior order- q -subplane with exterior splash $\mathbb{S}$. Consider the involutory homography Δ' of $\text{PG}(2, q^3)$ with matrix

$$\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

it fixes ℓ_∞ . As the homography Δ of ℓ_∞ given in Lemma 3.8 exchanges the two families $\mathcal{X}$, $\mathcal{Y}$, of order- q -sublines of $\mathbb{S}$, Δ' exchanges the two sets of order- q -sublines in $\mathbb{S}$, that is, Δ' maps $\mathcal{X}$ to $\mathcal{Y}$ and $\mathcal{Y}$ to $\mathcal{X}$. Thus Δ' does not fix π , as only elements of the Singer group $I = \text{PGL}(3, q^3)_{\pi, \ell_\infty}$ fix π and ℓ_∞ , and I acts regularly on the elements of $\mathcal{X}$ by Theorem 3.1. Thus Δ' fixes $\mathbb{S}$ and maps π to another exterior order- q -subplane π' . Hence Δ' maps the π -pencil-sublines to the π' -pencil-sublines. Thus if $\mathcal{X}$ contains the π -pencil-sublines, then $\mathcal{Y}$ contains the π' -pencil-sublines, as required. $\square$

Note that there is an analogous argument for scattered linear sets. Suppose that the exterior splash/scattered linear set $\mathbb{S}$ in the above proof is the projection of an order- q -subplane α from a point P onto ℓ_∞ , then under the homography Δ' , we obtain the same linear set $\mathbb{S}$ as the projection of an order- q -subplane α' from a point P' , and the order- q -sublines which are regular with respect to α are now irregular with respect to α' . That is, the notion of regular and irregular sublines in a fixed scattered linear set $\mathbb{S}$ of rank 3 can be interchanged by considering a different associated order- q -subplane that projects the linear set $\mathbb{S}$.

5 Projection of order- q -subplanes

In this section we look further at the relationship between exterior splashes and linear sets. Lunardon and Polverino [16] showed that a $\text{GF}(q)$ -linear set of rank 3 of $\text{PG}(1, q^3)$ can be constructed by projecting an order- q -subplane α of $\text{PG}(2, q^3)$ from a point $P \notin \alpha$ onto $\ell_\infty \cong \text{PG}(1, q^3)$. We compare this with our definition of the splash of an order- q -subplane π onto ℓ_∞ , namely the intersection of the lines of π with ℓ_∞ . If π is exterior to ℓ_∞ , then the exterior splash of π is equivalent to a $\text{GF}(q)$ -linear set of rank 3 and size $q^2 + q + 1$. If π is tangent to ℓ_∞ , then [3, Theorem 7.1] shows that the tangent splash of π is equivalent to a $\text{GF}(q)$ -linear set of rank 3 and size $q^2 + 1$. This leads to the question: given an order- q -subplane π , can the splash of π on ℓ_∞ be the same set of points as the projection of π onto ℓ_∞ from some point? We consider the case when π is exterior to ℓ_∞ in Section 5.1, and consider the case when π is tangent to ℓ_∞ in Section 5.2.

5.1 Projection and exterior splashes

In this section we explain under what circumstances the exterior splash of an exterior order- q -subplane π is equal to the projection of π onto ℓ_∞ . We also discuss a conjecture regarding which exterior splashes of ℓ_∞ can be obtained by projecting a fixed exterior plane order- q -subplane onto ℓ_∞ .

We begin by showing that the Singer cycle of Theorem 3.1 acts regularly on the special conics, and special-dual conics of an order- q -subplane.

Theorem 5.1 *Let π be an order- q -subplane of $\text{PG}(2, q^3)$ exterior to ℓ_∞ . The group $I = \text{PGL}(3, q^3)_{\pi, \ell_\infty}$ acts regularly on the set of (π, ℓ_∞) -special conics of π , and acts regularly on the (π, ℓ_∞) -special-dual conics of π .*

Proof By Theorem 3.1, I fixes π and hence acts on the projective plane $\mathcal{P}$ with points the points of π , and lines the set $\mathcal{C}$ of (π, ℓ_∞) -special conics of π . Consider the orbit $\mathcal{C}^I$ of $\mathcal{C}$ under I . As I is transitive on the points of π there is a constant number n of elements of $\mathcal{C}^I$ through each point of π . Count the pairs $(Q, \mathcal{D})$ where Q is a point of π on a (π, ℓ_∞) -special conic $\mathcal{D} \in \mathcal{C}^I$. We have

$$(q^2 + q + 1) \times n = |\mathcal{C}^I| \times (q + 1).$$

As $q^2 + q + 1$ and $q + 1$ have no common factors, it follows that $q + 1$ divides n . However, as $\mathcal{P}$ is a projective plane, $n \leq q + 1$. Hence $n = q + 1$ and so $|\mathcal{C}^I| = q^2 + q + 1$. Similarly, I acts regularly on the (π, ℓ_∞) -special-dual conics of π . $\square$

Theorem 5.2 *Let π be an exterior order- q -subplane with exterior splash $\mathbb{S}$ on ℓ_∞ , carriers E_1, E_2 , and third conjugate point E_3 . Let P be a point of $\text{PG}(2, q^3) \setminus \pi$, then the projection of π from P onto ℓ_∞ is equal to $\mathbb{S}$ if and only if q is even, and $P = E_3$.*

Proof By Theorem 3.3, we can without loss of generality prove this for the order- q -subplane $\pi = \text{PG}(2, q)$ with exterior splash $\mathbb{S}$ onto the exterior line $\ell = [-\tau\tau^q, \tau^q + \tau, -1]$. By Theorem 3.1, the group $I = \text{PGL}(3, q^3)_{\pi, \ell}$ fixing π and ℓ is cyclic of order $q^2 + q + 1$; acts regularly on the points and lines of π ; and by Theorem 5.1 acts regularly on the (π, ℓ) -special-dual conics of π . By Theorem 4.2, there are $2(q^2 + q + 1)$ order- q -sublines in $\mathbb{S}$, divided into two families denoted $\mathcal{X}$ and $\mathcal{Y}$, one family contains the π -pencil-sublines of $\mathbb{S}$, and the other family contains the π -dual-conic-sublines. So I acts regularly on the order- q -sublines of $\mathcal{X}$, and acts regularly on the order- q -sublines of $\mathcal{Y}$.

Let P be a point of $\text{PG}(2, q^3)$ not in π or ℓ , and let L_P be the projection of π onto ℓ from P . Suppose that $L_P = \mathbb{S}$, that is, the projection of π from P is the same as the exterior splash of π onto ℓ . By [12], the $q^2 + q + 1$ order- q -sublines of π are projected onto order- q -sublines of $L_P = \mathbb{S}$ that lie in the same family, $\mathcal{X}$ say. Further, the line joining P to an order- q -subline d in the other family $\mathcal{Y}$ meets π in a set of points which form a conic of π whose extension to $\text{PG}(2, q^3)$ contains P , and hence P^q and P^{q^2} . That is, it is a (π, PP^q) -special conic, and so

the $q^2 + q + 1$ (π, PP^q) -special conics of π are projected to the $q^2 + q + 1$ order- q -sublines in $\mathcal{Y}$.

As the group I acts regularly on the order- q -sublines of $\mathcal{Y}$, it induces a Singer cycle on the $q^2 + q + 1$ (π, PP^q) -special conics of π . All these conics have P, P^q, P^{q^2} in common. If P is not a fixed point of this Singer cycle, then the $q^2 + q + 1$ (π, PP^q) -special conics all have the images of P in common, a contradiction as two distinct irreducible conics have at most four points in common. Thus P is a fixed point of the Singer cycle, and $P \notin \ell$, so $P = \ell^q \cap \ell^{q^2}$. That is, if $L_P = \mathbb{S}$ then $P = \ell^q \cap \ell^{q^2} = (1, \tau^{q^2}, \tau^{2q^2})$.

It remains to show that for the point $P = \ell^q \cap \ell^{q^2}$, $L_P = \mathbb{S}$ if and only if q is even. By Example 3.2, we can parameterise the points of $\mathbb{S}$ as $E + \theta E^q$ for $\theta \in \text{GF}(q^3)$, $\theta^{q^2+q+1} = -1$. Let (x, y, z) , $x, y, z \in \text{GF}(q)$, be a point of π . The line joining this point to $P = (r, s, t)$ has coordinates $[sz - ty, tx - rz, ry - sx]$. It meets the line $\ell = [-\tau\tau^q, \tau^q + \tau, -1]$ in the point $E + \theta E^q$ with parameter

$$\theta = \frac{x(s\tau^2 - t\tau) + y(t - r\tau^2) + z(r\tau - s)}{x(t\tau^q - s\tau^{2q}) + y(r\tau^{2q} - t) + z(s - r\tau^q)} = \frac{f(x, y, z)}{g(x, y, z)}.$$

If $P = (r, s, t) = (1, \tau^{q^2}, \tau^{2q^2})$ we see that $f^{q^2} = g$, and so $\theta^{q^2+q+1} = 1$. Now $\mathbb{S}$ is the set of points with parameter θ , where $\theta^{q^2+q+1} = -1$, hence the projection L_P from $P = (1, \tau^{q^2}, \tau^{2q^2})$ is the same as the exterior splash $\mathbb{S}$ of π onto ℓ if and only if q is even. $\square$

Suppose we fix an order- q -subplane π and an exterior line ℓ . We consider the set $\mathcal{S}$ of all the projections of π from a point $P \in \text{PG}(2, q^3) \setminus \{\pi, \ell\}$ onto ℓ . Note that the projections in $\mathcal{S}$ are either tangent or exterior splashes. As there are less than q^6 possible points to project π from, and more than q^6 exterior splashes, we will not obtain all the possible exterior splashes on ℓ . We further examine this situation. As usual, let $I = \text{PGL}(3, q^3)_{\pi, \ell}$ with fixed points E_1, E_2, E_3 and fixed lines ℓ, m, n . Let J be the subgroup of $\text{PGL}(3, q^3)$ that fixes E_1, E_2, E_3 . Then it is straightforward to show that I is a normal subgroup of J , and that further J has seven orbits, namely E_1 ; E_2 ; E_3 ; $\ell \setminus \{E_1, E_2\}$; $m \setminus \{E_1, E_3\}$; $n \setminus \{E_2, E_3\}$; and the points $\text{PG}(2, q^3) \setminus \{\ell, m, n\}$. Moreover, the orbits of I on ℓ are E_1, E_2 and $q - 1$ exterior splashes, with similar orbits on m and n . Finally the orbits of I on $\text{PG}(2, q^3) \setminus \{\ell, m, n\}$ consist of order- q -subplanes. We now look at the action of I on the splashes in $\mathcal{S}$.

Lemma 5.3 *1. An exterior splash in $\mathcal{S}$ is either fixed under I , or has an orbit of size $q^2 + q + 1$ under I .
2. A tangent splash in $\mathcal{S}$ has an orbit of size $q^2 + q + 1$ under I .*

Proof For part 1, consider an exterior splash $\mathbb{S} \in \mathcal{S}$ with an orbit under I of size less than $|I| = q^2 + q + 1$, so $I_{\mathbb{S}}$ is non-trivial. We show that in this case $\mathbb{S}$ is fixed by I . Let $g \in I_{\mathbb{S}}$, g not the identity, so g fixes E_1 and E_2 . As I is cyclic, it is abelian, so $\langle g \rangle$ is a normal subgroup of I . Thus $\langle g \rangle$ fixes either all the points in an orbit of I , or has no fixed points on an orbit of I . In the first case, g fixes at least $(q^2 + q + 1) + 2$ points of ℓ , which is more points than an order- q -subline of ℓ contains, hence g fixes ℓ pointwise, and so g is the identity on ℓ . As I is

faithful on ℓ , it follows that g is the identity. Thus the second case occurs. From Lemma 3.8 we can show that the full group (acting faithfully) on the splash is of size $2(q^2 + q + 1)$, consisting of $q^2 + q + 1$ involutions and the cyclic group I of odd order, whose elements only fix the carriers of the splash. As $g \in I_{\mathbb{S}} \subset I$, it follows that g has odd order and the only fixed points are the carriers of $\mathbb{S}$. This implies that $\mathbb{S}$ has carriers E_1, E_2 , and is fixed by I . So an exterior splash in $\mathcal{S}$ has orbit of size 1 or $|I| = q^2 + q + 1$.

For part 2, as the exterior splash of π on ℓ does not contain E_1 or E_2 , by definition no line of π contains E_1 or E_2 , thus the tangent splash obtained by projecting π from a point on a line of π does not have centre E_1 or E_2 . Thus the centre of the tangent splash is not a fixed point of I , hence its orbit under I is of size $|I|$, and hence the $q^2 + q + 1$ tangent splash projections in $\mathcal{S}$ have different centres, and so are distinct. $\square$

Lemma 5.4 *Let $\mathbb{S}$ be an exterior splash in $\mathcal{S}$ with carriers E_1, E_2 . Then if q is even, $\mathbb{S}$ is the projection of π from 1 point or $q^2 + q + 1$ points; and if q is odd, $\mathbb{S}$ is the projection of π from $q^2 + q + 1$ or $q^2 + q + 2$ points.*

Proof We can without loss of generality consider the order- q -subplane $\pi = \text{PG}(2, q)$ and the exterior line $\ell = [-\tau\tau^q, \tau^q + \tau, -1]$. First note that the projection of π from E_3 is an exterior splash with carriers E_1, E_2 . As $\mathbb{S}$ occurs as a projection of π , either it is from E_3 (this would give the 1 point or the $q^2 + q + 2$ points case), or it is from a point P belonging to an orbit θ under I of size $q^2 + q + 1$. In this case, as $\mathbb{S}$ and π are both fixed by I , it follows that every point of the orbit θ projects π on to the same splash $\mathbb{S}$. Now suppose there is another point W which also projects π onto $\mathbb{S}$. Recall that the exterior splash $\mathbb{S}$ has two families of order- q -sublines, $\mathcal{X}$ and $\mathcal{Y}$. An order- q -subline m of π is projected by P onto an order- q -subline in one of these families, $\mathcal{X}$ say. Further, the images of m under I are projected onto order- q -sublines in $\mathcal{X}$. Thus W cannot project m onto an order- q -subline of $\mathcal{X}$, since the point of projection of an order- q -subline is unique. Thus W projects m onto an order- q -subline y of the family $\mathcal{Y}$. Similar to the proof of Theorem 5.2, the lines joining W to the points of y meet π in a set of points which form a conic of π whose extension to $\text{PG}(2, q^3)$ contains W , and hence W^q and W^{q^2} . Then, similar to the proof of Theorem 5.2, it follows that W is a fixed point of the Singer cycle I , and so $W = \ell^q \cap \ell^{q^2}$. By Theorem 5.2, if q is odd, then there is at most one exterior splash with carriers E_1, E_2 with $q^2 + q + 2$ points of projection, and if q is even, all the exterior splashes with carriers E_1, E_2 have at most $q^2 + q + 1$ points of projection. $\square$

We conject that the following fully describes the projection of an order- q -subplane onto an exterior line.

Conjecture 5.5 *Let π be an order- q -subplane in $\text{PG}(2, q^3)$ with exterior line ℓ . Let $\mathcal{S}$ be the set of all the projections of π from a point $P \in \text{PG}(2, q^3) \setminus \{\pi, \ell\}$ onto ℓ . Note that the projections in $\mathcal{S}$ are either tangent or exterior splashes. More specifically,*

1. *The points $P \in \text{PG}(2, q^3) \setminus \{\pi, \ell\}$ that lie on the extension of a line of π project π onto distinct tangent splashes. Hence there are $(q^2 + q + 1)(q^3 - q - 1)$ such tangent splashes in $\mathcal{S}$.*

2. The exterior splashes in $\mathcal{S}$ can be divided into three distinct groups. Let $\mathbb{S}_0$ be the exterior splash of π onto ℓ , and let $\mathbb{S}_1, \dots, \mathbb{S}_{q-2}$ be the distinct exterior splashes on ℓ that have the same carriers E_1, E_2 as $\mathbb{S}_0$.
- (a) i. If q is even, then $\mathbb{S}_0$ is a projection of π from E_3 , and $\mathbb{S}_i, i = 1, \dots, q-2$ is a projection from exactly $q^2 + q + 1$ points which lie in one orbit of I .
 - ii. If q is odd, then $\mathbb{S}_i, i = 1, \dots, q-2$ is the projection of π from exactly $q^2 + q + 1$ points which lie in one orbit of I , with the exception of one of these which is also a projection of π from E_3 .
 - (b) The remaining exterior splashes in $\mathcal{S}$ are either:
 - i. projections of π from exactly two points which are not in the same orbit under I ; or
 - ii. projections of π from exactly one point.

5.2 Projection and tangent splashes

Let π be an order- q -subplane that is tangent to ℓ_∞ at the point T , that is, π is a tangent order- q -subplane. The lines of π meet ℓ_∞ in a set $\mathcal{S}_T$ of $q^2 + 1$ points (including T) called a *tangent splash* with *centre* T . For a detailed study of tangent splashes, see [3], in particular, by [3, Theorem 7.1], tangent splashes are projectively equivalent to $\text{GF}(q)$ -linear sets of rank 3 and size $q^2 + 1$. We can construct such a linear set by projecting an order- q -subplane. We show that the linear set obtained by projecting a tangent order- q -subplane π onto ℓ_∞ can never equal the tangent splash of π onto ℓ_∞ .

Lemma 5.6 *Let ℓ, m be lines of $\text{PG}(2, q^3)$, and b an order- q -subline of m disjoint from ℓ . Then each order- q -subline of ℓ , disjoint from m , is the projection of b from exactly one point P not on ℓ or m .*

Proof We first show that the subgroup K of $\text{PGL}(2, q^3)$ (acting on $\text{PG}(1, q^3)$) fixing a point P of $\text{PG}(1, q^3)$ is regular on the order- q -sublines not containing P . Note that $\text{PGL}(2, q^3)$ acts faithfully on an order- q -subline b of $\text{PG}(1, q^3)$, as any such homography fixing 3 points is the identity. Thus $|\text{PGL}(2, q^3)_b| = |\text{PGL}(1, q)| = q(q^2 - 1)$. As $\text{PGL}(2, q^3)$ is sharply 3-transitive on the points of $\text{PG}(1, q^3)$, it is transitive on the order- q -sublines of $\text{PG}(1, q^3)$. Consider the subline $b = \text{PG}(1, q)$ of $\text{PG}(1, q^3)$. From above, the subgroup H of $\text{PGL}(2, q^3)$ fixing b is of order $q(q^2 - 1)$. If we consider a homography acting on $P = (\tau, 1)^t$, a straightforward calculation shows that H_P is just the identity, hence by the orbit stabilizer theorem, H is transitive on the points of $\text{PG}(1, q^3)$ not on b . Now suppose P is a point not on b . If there is an element of $K = \text{PGL}(2, q^3)_P$ which fixes b , then this element belongs to H and fixes the point P outside b , and so is the identity. Thus K acts semiregularly on the $q^3(q^3 - 1)$ order- q -sublines not through P , but as $|K| = q^3(q^3 - 1)$, this action is transitive, and hence regular.

Consider the group L of axial homographies with axis m and centre on ℓ . There are $q^3 - 1$ non-identity elations in L , and $q^3(q^3 - 2)$ non-identity homologies in L , so $|L| = q^3(q^3 - 1)$.

As an element of L fixes ℓ and $P = \ell \cap m$, it induces a homography σ on ℓ which fixes P . As the subgroup K above is regular on the order- q -sublines not containing P , σ is either the identity, or acts semi-regularly on the order- q -sublines of ℓ not through P . If σ acts as the identity on ℓ , then as it has axis m , it follows that it is the identity on $\text{PG}(2, q^3)$. Hence L acts semi-regularly and hence regularly on the order- q -sublines of ℓ not through P .

Now consider a point X projecting b onto an order- q -subline c on ℓ . Under the action of L , b is fixed pointwise (as b is contained in the axis of the elements of L) and c is mapped to the $q^3(q^3 - 1)$ order- q -sublines on ℓ not through P . Further, as L acts semi-regularly on the $q^3(q^3 - 1)$ points of $\text{PG}(2, q^3)$ not on $\ell \cup m$, it follows that L is regular on the points not on $\ell \cup m$. Thus distinct points of $\text{PG}(2, q^3)$ not on $\ell \cup m$ project b onto distinct order- q -sublines of ℓ . $\square$

Theorem 5.7 *Let π be a tangent order- q -subplane with tangent splash $\mathcal{S}_T$ on ℓ_∞ . Then $\mathcal{S}_T$ is not the projection of π onto ℓ_∞ from a point P .*

Proof We let $L_{X,P}$ denote the projection of the set of points X of $\text{PG}(2, q^3)$ onto ℓ_∞ from the point P . Note that if we project the tangent order- q -subplane π onto ℓ_∞ from a point $P \in \ell_\infty$ or $P \in \pi$, then $L_{\pi,P}$ does not have enough points to be the tangent splash $\mathcal{S}_T$ of π . Similarly, if P is a point not on a line of π , then $L_{\pi,P}$ has too many points to be a tangent splash. Suppose we project from a point $P \notin \pi$ that is on a line ℓ of π not through the centre T of $\mathcal{S}_T$, and suppose $L_{\pi,P} = \mathcal{S}_T$. Let b be an order- q -subline of π not through T , and not on ℓ , then $L_{b,P}$ is an order- q -subline of $\mathcal{S}_T$ not through the centre T , contradicting [3, Corollary 7.3]. So this case cannot occur.

So, suppose we project π from a point $P \notin \pi$ that lies on a line of π through the centre T , and suppose $L_{\pi,P} = \mathcal{S}_T$. Let J be the subgroup of $\text{PGL}(3, q^3)_\pi \cong \text{PGL}(3, q)$ that fixes T and ℓ_∞ . Consider J_P , the subgroup of J fixing the projection point P . From the proof of [3, Theorem 4.2], J contains only central collineations with centre T and axes belonging to π . So P is fixed by a non-identity element σ of J only if it lies on the axis of σ . So J_P is the set of q elations with centre T and axis TP . Hence $|P^J| = |J|/|J_P| = q(q - 1)$.

Let b be an order- q -subline of π not through T . Consider the projection $c = L_{b,P}$. If Q is another point in P^J , then by Lemma 5.6, it follows that $L_{b,Q}$ is not c . Thus the projections of b under the $q(q - 1)$ different points in P^J is the orbit of c under J , and is of size $q(q - 1)$.

However, each point A of π determines a unique order- q -subline of $\mathcal{S}_T$ via the intersection of the lines of π through A with ℓ_∞ , and all the order- q -sublines in the tangent splash $\mathcal{S}_T$ are determined this way (by [3, Corollary 7.4]). As the points of π under J fall into $q + 1$ orbits, each consisting of q points lying on a line through T , it follows that $|c^J| = q$, contradicting the previous paragraph. $\square$

6 Order- q -subplanes with a common exterior splash

In this section we investigate the intersection of two exterior order- q -subplanes that have a common exterior splash. We begin with a counting result.

Theorem 6.1 *Given an exterior splash $\mathbb{S}$ on ℓ_∞ , there are $2q^6(q^3 - 1)$ order- q -subplanes with exterior splash $\mathbb{S}$.*

Proof By Theorem 3.5, the number of exterior splashes is equal to the number of covers of a circle geometry $CG(3, q)$, which is $\frac{1}{2}q^3(q^3 + 1)(q - 1)$. Counting quadrangles gives that there are $q^9(q^3 - 1)(q^3 + 1)(q - 1)$ order- q -subplanes exterior to a given line ℓ_∞ . Further, by Theorem 3.3, the group of homographies of $PG(2, q^3)$ fixing ℓ_∞ is transitive on the order- q -subplanes exterior to ℓ_∞ . Hence the number of order- q -subplanes with a fixed exterior splash is the total number of order- q -subplanes exterior to ℓ divided by the number of exterior splashes on ℓ , that is, $2q^6(q^3 - 1)$. $\square$

Theorem 6.2 *Let $\mathbb{S}$ be an exterior splash of ℓ_∞ and let ℓ be an order- q -subline exterior to ℓ_∞ , whose extension to $PG(2, q^3)$ contains a point of $\mathbb{S}$. Then there are exactly two order- q -subplanes that contain ℓ and have exterior splash $\mathbb{S}$.*

Proof Let π be an order- q -subplane with exterior splash $\mathbb{S}$, and let ℓ be a line of π . Consider the subgroup $I = G_{\pi, \ell_\infty}$ of $G = PGL(3, q^3)$ fixing π and ℓ_∞ . By Theorem 3.1 the group I fixes $\mathbb{S}$ and is transitive on the points of $\mathbb{S}$. By [4, Theorem 2.5], the subgroup of G fixing ℓ point-wise is transitive on the exterior order- q -sublines on a line $m \neq \ell$. Thus, given an exterior splash $\mathbb{S}$, the group fixing $\mathbb{S}$ is transitive on the exterior order- q -sublines lying on a line through a point of $\mathbb{S}$. Hence the number n of exterior order- q -subplanes containing an order- q -subline ℓ lying on a line through a point of a given exterior splash $\mathbb{S}$ is a constant. Thus if we count the pairs (π, ℓ) where π is an exterior order- q -subplane with a given exterior splash $\mathbb{S}$, containing any of the $q^3(q^3 - 1)$ order- q -sublines ℓ lying on any of the q^3 lines through a point of a given exterior splash $\mathbb{S}$, using Theorem 6.1, we get

$$2q^6(q^3 - 1) \times (q^2 + q + 1) = (q^2 + q + 1) \times q^3 \times q^3(q^3 - 1) \times n$$

and so $n = 2$ as required. $\square$

Recall that three collinear points lie in a unique order- q -subline, and a quadrangle lies in a unique order- q -subplane. Hence two order- q -subplanes in $PG(2, q^3)$ can meet in either 0, 1, 2 or 3 points, or in $q + 1$ points lying on an order- q -subline, or in $q + 2$ points containing an order- q -subline. In Lemma 6.2, we showed that there are exactly two exterior order- q -subplanes that share an order- q -subline and have the same exterior splash. We now show that these two order- q -subplanes do not have any further common points.

Theorem 6.3 *Let π_1, π_2 be two exterior order- q -subplanes with a common exterior splash $\mathbb{S}$ on ℓ_∞ . Then π_1, π_2 meet in at most $q + 1$ points, that is, π_1, π_2 meet in 0, 1, 2 or 3 points, or in an order- q -subline.*

Proof By Theorem 3.3 and Example 3.2, we can without loss of generality let $\pi_1 = \text{PG}(2, q)$ with exterior splash $\mathbb{S}_1 = \{E_1 + \theta E_2 : \theta^{q^2+q+1} = -1, \theta \in \text{GF}(q^3)\}$ on the exterior line $\ell = [-\tau\tau^q, \tau^q + \tau, -1]$ where $E_1 = (1, \tau, \tau^2)$, $E_2 = E_1^q = (1, \tau^q, \tau^{2q})$ are the carriers of $\mathbb{S}_1$. Let π_2 be an order- q -subplane exterior to ℓ with exterior splash $\mathbb{S}_2$ on ℓ . We show that if $\mathbb{S}_1 = \mathbb{S}_2$ and π_1, π_2 contain $q + 2$ common points, then $\pi_1 = \pi_2$. By Theorem 3.1, as the group I fixing π_1 and ℓ (and hence $\mathbb{S}_1$) is transitive on the lines of π_1 , without loss of generality we can suppose π_2 contains the order- q -subline $m = \{M_b = (0, 1, b) : b \in \text{GF}(q) \cup \{\infty\}\}$ of π_1 . Suppose that π_2 and π_1 contain a further common point P not on m , so $P = (\mu, \nu, \omega)$ with $\mu, \nu, \omega \in \text{GF}(q^3)$, with $\mu \neq 0$. As $P \in \pi_1$, we have $\nu/\mu, \omega/\mu \in \text{GF}(q)$.

Let $Q = (0, 0, 1) = M_\infty \in m$, so the line PQ contains an order- q -subline of π_2 . Without loss of generality, by suitable choice of $\mu, \nu, \omega \in \text{GF}(q^3)$, let that order- q -subline be determined by the three points P , Q and $T = P + Q = (\mu, \nu, \omega + 1)$. That is, π_2 is determined by the quadrangle $(0, 1, 1)$, $(0, 1, 0)$, $P = (\mu, \nu, \omega)$ and $T = (\mu, \nu, \omega + 1)$. Further, π_1 and π_2 both contain the order- q -subline m and the point P .

We will show that, if $\mathbb{S}_2 = \mathbb{S}_1$, then $\mu \in \text{GF}(q)$, and hence $\nu, \omega \in \text{GF}(q)$ and so $T \in \pi_1$. The order- q -subline n defined by $P, Q, T = P + Q$ has points $N_c = P + cQ = (\mu, \nu, \omega + c)$ for $c \in \text{GF}(q)$. As the points M_b, N_c lie in π_2 for $b, c \in \text{GF}(q)$, the line M_bN_c meets ℓ in a point X_{bc} of $\mathbb{S}_2$. Now M_bN_c has coordinates $[c - b\nu + \omega, b\mu, -\mu]$, and it meets ℓ in the point $X_{bc} = E_1 + \theta_{bc}E_2$ with

$$\theta_{bc} = -\frac{c + b(\mu\tau - \nu) - (\mu\tau^2 - \omega)}{c + b(\mu\tau^q - \nu) - (\mu\tau^{2q} - \omega)}.$$

The point X_{bc} is in the exterior splash $\mathbb{S}_1$ if and only if $\theta_{bc}^{q^2+q+1} = -1$. We can regard this equation as a multivariate polynomial of degree 3 in b and c . Note that if a multivariate polynomial of degree $k < q$ over $\text{GF}(q^3)$ is identically zero over $\text{GF}(q)$, then the polynomial is the zero polynomial. If $q = 2$ or 3 , a computer search using Magma [6] verifies the result holds, so we assume $q > 3$. Expanding, and calculating the coefficients of bc^2 and c^2 , and equating them to zero gives

$$\begin{aligned} 0 &= \kappa\tau + \kappa^q\tau^q + \kappa^{q^2}\tau^{q^2} \\ 0 &= \kappa\tau^2 + \kappa^q\tau^{2q} + \kappa^{q^2}\tau^{2q^2}, \end{aligned}$$

respectively, where $\kappa = \mu - \mu^{q^2}$, and so $\kappa + \kappa^q + \kappa^{q^2} = 0$. That is, the point $M_bN_c \cap \ell$ lies in the exterior splash $\mathbb{S}_1$ for all $b, c \in \text{GF}(q)$ if

$$\begin{pmatrix} 1 & 1 & 1 \\ \tau & \tau^q & \tau^{q^2} \\ \tau^2 & \tau^{2q} & \tau^{2q^2} \end{pmatrix} \begin{pmatrix} \kappa \\ \kappa^q \\ \kappa^{q^2} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}.$$

This 3×3 matrix is invertible, so the only solution is $\kappa = \kappa^q = \kappa^{q^2} = 0$, that is $\mu = \mu^{q^2}$, and so $\mu \in \text{GF}(q)$. Hence π_2 has exterior splash $\mathbb{S}_1$ if and only if $\mu \in \text{GF}(q)$. As $\nu/\mu, \omega/\mu \in \text{GF}(q)$, it follows that $T \in \pi_2 \cap \pi_1$. Thus π_2 and π_1 share the quadrangle $P, T, (0, 1, 0), (0, 1, 1)$ and so $\pi_2 = \pi_1$ as required. $\square$

Suppose π_1, π_2 are two exterior order- q -subplanes with a common exterior splash $\mathbb{S}$ that meet in $q + 1$ points. We conject that the two families of order- q -sublines of $\mathbb{S}$ with respect to π_1, π_2

are *swapped*. That is, π_1, π_2 correspond to the order- q -subplanes π, π' with order- q -sublines behaving as in Theorem 4.4.

7 Conclusion

In this article we looked at the exterior splash as a set of points of $\text{PG}(1, q^3)$, and showed its equivalence to $\text{GF}(q)$ -linear sets of rank 3 and size $q^2 + q + 1$, Sherk surfaces of size $q^2 + q + 1$, and covers of the circle geometry $CG(3, q)$. We also investigated properties of an exterior order- q -subplane and its corresponding exterior splash, in particular relating order- q -sublines of an exterior splash to the corresponding order- q -subplane. Further, we looked at how our construction of exterior splashes related to the projection construction of linear sets.

In future work, we look in the Bruck-Bose representation of $\text{PG}(2, q^3)$ in $\text{PG}(6, q)$, and study exterior splashes in this context, further utilising properties of scattered linear sets of rank 3, as well as properties of hyper-reguli in $\text{PG}(5, q)$.

References

- [1] R.D. Baker, J.M.N. Brown, G.L. Ebert and J.C. Fisher. Projective bundles. *Bull. Belg. Math. Soc.*, **3** (1994) 329–336.
- [2] S.G. Barwick and W.-A. Jackson. A characterisation of tangent subplanes of $\text{PG}(2, q^3)$. *Des. Codes Cryptogr.*, **71** (2014) 541–545.
- [3] S.G. Barwick and W.-A. Jackson. An investigation of the tangent splash of a subplane of $\text{PG}(2, q^3)$. *Des. Codes Cryptogr.*, **76** (2015) 451–468.
- [4] S.G. Barwick and W.-A. Jackson. The tangent splash in $\text{PG}(6, q)$. *Discrete Math.*, **338** (2015) 1178–1190.
- [5] A. Blokhuis and M. Lavrauw. Scattered spaces with respect to a spread in $\text{PG}(n, q)$. *Geom. Dedicata*, **81** (2000) 231–243.
- [6] W. Bosma, J. Cannon, and C. Playoust, The Magma algebra system. I. The user language, *J. Symbolic Comput.*, 24 (1997), 235–265.
- [7] R.H. Bruck. Construction problems of finite projective planes. *Conference on Combinatorial Mathematics and its Applications*, University of North Carolina Press, (1969) 426–514.
- [8] R.H. Bruck. Circle geometry in higher dimensions. *A Survey of Combinatorial Theory*, eds. J. N. Srivastava et al, Amsterdam (1973) 69–77.
- [9] R.H. Bruck. Circle geometry in higher dimensions. II. *Geom. Dedicata*, **2** (1973) 133–188.

- [10] C. Culbert and G.L. Ebert. Circle Geometry and three-dimensional subregular translation planes. *Innov. Incidence Geom.*, **1** (2005) 3–18.
- [11] G. Donati and N. Durante. Scattered linear sets generated by collineations between pencils of lines, *J. Algebraic. Comb.*, **40** (2014) 1121–1134.
- [12] M. Lavrauw and G. Van de Voorde. On linear sets on a projective line. *Des. Codes Cryptogr.*, **56** (2010) 89–104.
- [13] M. Lavrauw and G. Van de Voorde. Scattered linear sets and pseudoreguli. *Electronic J. Combin.*, **20** (2013).
- [14] M. Lavrauw and C. Zanella. Subgeometries and linear sets on a projective line. *Finite Fields Appl.*, to appear.
- [15] G. Lunardon, G. Marino, O. Polverino and R. Trombetti. Maximum scattered linear sets of pseudoregulus type and the Segre variety $\mathcal{S}_{n,n}$. *J. Algebraic. Comb.*, **39** (2014) 807–831.
- [16] G. Lunardon and O. Polverino. Translation ovoids of orthogonal polar spaces. *Forum Math.*, **16** (2004) 663–669.
- [17] F.A. Sherk. The Geometry of $\text{GF}(q^3)$. *Canad J. Math.*, **38** (1986) 672–696.